atplab

Medicinal Ingredients/Ingrédients médicinaux

per softgel/ par gélule

Borage Oil/Huile de bourrache (Borago officinalis)	120 mg
CLA (Conjugated Linoleic Acid)/ALC (Acide Linoléique conjugué)	450 mg
Fish Oil (sardines, anchovies, mackerels)/ Huile de poisson (sardines, anchois, maquereaux)	450 mg
Providing/Fournissant : EPA	130 mg
DHA	100 mg
Oleuropein 20 %/Oleuropéine 20 %	60 mg

Non-medicinal Ingredients: Beeswax, coconut oil, sunflower lecithin, gelatin, glycerin, carob extract. **Recommended Use or Purpose:** Source of omega-3 fatty acids for the maintenance of good health. Helps support cognitive health and/or brain function. May help to support a modest reduction in fat mass when used with a program of reduced intake of dietary calories and increased physical activity. **Recommended Dose:** Adults: 4 softgels/twice daily. Take with food. **Duration of use:** Consult a healthcare practitioner for use beyond 6 months. **Warnings:** Consult a healthcare practitioner prior to use if you are obese or have cardiovascular desease (CVD) risk factors (high blood pressure, high cholesterol and/or triglycerides), if you are pregnant or breastfeeding, or if your goal is to achieve weight loss. **Contra-indications :** Do not use if you have CVD, diabetes, metabolic syndrome or insulin resistance. **Known Adverse Reactions:** Some people may experience gastrointestinal upset.

Ingrédients non médicinaux : Cire d'abeille, huile de noix de coco, lécithine de tournesol, gélatine, glycérine, extrait de caroube. **Usage ou fins recommendés :** Source d'acides gras oméga-3 pour le maintien d'une bonne santé. Aide à contribuer à la santé cognitive et/ou aux fonctions cérébrales. Pourrait contribuer à une légère diminution de la masse adipeuse, dans le cadre d'un programme qui recommande une réduction de l'apport calorique et une augmentation de l'activité physique. **Dose recommandée :** Adultes : 4 gélules/2 fois par jour. Prendre avec de la nourriture. **Durée d'utilisation :** Consultez un médecin si l'usage se prolonge au-delà de 6 mois. **Mises en garde :** Consultez un médecin avant d'en faire l'usage si votre but est de perdre du poids, si vous êtes enceinte ou si vous allaitez, si vous êtes obèse ou si vous avez des facteurs de risque de maladies cardiovasculaires (MCV) (hypertension artérielle, taux élevé de cholestérol et/ou de triglycérides). **Contre-indication :** Ne pas utiliser si vous souffrez d'une MCV, de diabète, du syndrome métabolique ou si vous avez une résistance à l'insuline. **Effets indésirables connus :** Certaines personnes pourraient souffrir de troubles gastrointestinaux.

CONTACT US

1 888 770-0079

info@atplab.com